arXiv:0803.0981v1 [hep-th] 6 Mar 2008

Free-fall in a uniform gravitational field in non-commutative quantum mechanics: on an upper-bound on θ and the "quantum Galileo experiment"

K. H. C. Castello-Branco [a†] *and A. G. Martins* [b∗]

[a] *Instituto de Física de São Carlos, Universidade de São Paulo,*
Av. Trabalhador São-Carlense, 400. C.P. 369, 13560-970. São Carlos-SP,
Brazil

[b] *Instituto de Física, Universidade de Brasília,*
C. P. 04455, 70919-970. Brasília-DF, Brazil

Abstract

We study the gravitational quantum well in the context of Non-Commutative Quantum Mechanics (NCQM). Assuming non-commutativity of the canonical type between the coordinates of a two dimensional configuration space, we consider a neutral particle trapped in a gravitational well and *exactly* solve its energy eigenvalue problem. By resorting to experimental data from the *Grenoble experiment*, where the first energy levels of freely falling quantum ultracold neutrons were determined, we impose an upper-bound on the non-commutativity parameter, θ. We also investigate the time of flight of a quantum particle moving in a uniform gravitational field in NCQM. This is related to the Equivalence Principle. As we consider stationary, energy eigenstates, *i.e.*, delocalized states, the time of flight must be measured by a *quantum clock*, suitably coupled to the particle. By considering the clock as a small perturbation we solve the (stationary) scattering problem associated and show that the time of flight is equal to the classical result, when the measurement is made far from the turning point. This result is interpreted as an extension of the Equivalence Principle to the realm of NCQM.

† karlucio@ursa.ifsc.usp.br
∗ andrey@fis.unb.br

1 Introduction

The idea that space-time would have a *non-commutative* structure was proposed by Heisenberg and others [1], already at the very beginning of Quantum Field Theory (QFT), so that an effective ultraviolet cuttoff could be introduced at very small length scales, in an attempt to get rid of divergences. It was only in 1947, with Snyder [2], that this idea of space-time non-commutativity was formalized (for a historical introduction, see [3, 4]). But due to the success achieved by the renormalization program of QFTs, relatively low interest remained on the subject. In the late 1990s some results coming from String Theory have suggested that space-time may display a non-commutative structure [5], thus starting a great revival on the study of QFTs based on non-commuting space-time coordinates (for reviews, see [3, 6, 7]). Nevertheless, we remark that the issue of space-time non-commutativity was considered in an earlier work by Doplicher *et al.*, who constructed a unitary QFT based on a non-commutative space-time [8], motivated by the issue of the description of the quantum nature of the space-time.

As quantum mechanics can be considered the one-particle sector of quantum field theory, it is interesting to study the quantum mechanics defined on non-commutative spaces. Lately, this *Non-Commutative Quantum Mechanics* (NCQM) has been increasingly studied (see, for instance, Ref.'s [9]) and effects of non-commutativity that might be experimentally detectable have been investigated [10]. There have been also studies of many-particle systems [11] and an approach to NCQM that is directly related to the ideas developed in [8] has been developed in [12], by means of an algebraic setup in which the time is a non-commutative coordinate as well as the space coordinates, with interesting ensuing consequences (see Ref.'s [13, 14]).

In this paper we are concerned about the physics of a quantum particle in a uniform gravitational field. In the first part of the paper we study the problem known as the *gravitational quantum well.* This can be obtained by the Earth's gravitational field and a perfectly reflecting mirror at the bottom. In the context of ordinary quantum mechanics this well-known problem has been thoroughly studied in text-books and pedagogic articles (see, for instance, [15]). But this simple theoretical investigation gained a lot in importance, as recently such a quantum well was experimentally realized and the first two quantum states of neutrons moving in the Earth's gravitational well were identified. This was achieved through the *Grenoble experiment*, perfomed by Nesvizhevsky *et al.* [16]-[18]. In this paper we study the non-commutative version of this problem, by means of the solution of the non-commutative Schrödinger equation. As it will be shown, the experimental results obtained in [16]-[18] allow us to find an upper-bound on the parameter of the spatial non-commutativity. The first study to treat the non-commutative gravitational quantum well was done by Bertolami *et al.* [19], but in a non-commutative model different from the one we will consider in this paper.

We remark that we have assumed only *spatial* non-commutativity, in contrast to other works in the literature, which usually considered non-commutativity of both configuration and momentum spaces (see [19]-[22]). Those papers treated the non-commutative gravitational quantum well and used data of the Grenoble experiment [16]-[18] to find upper-bounds on the value of the *momentum-momentum* non-commutativity parameter. Nonetheless, differently from [19]-[22], we find the shifts in the energy levels due to spatial non-commutativity only. Moreover, we treat the problem *exactly*, without resorting to any perturbative approach.

In the second part of this paper we consider the issue of the *Equivalence Principle* in the context of NCQM. Already in ordinary quantum mechanics, the validity of the Equivalence Principle is an interesting and subtle question. The Equivalence Principle is generally claimed to be the root of the problem of fully conciliate Quantum Mechanics with General Relativity.

We address the issue of the Equivalence Principle in NCQM motivated by the study due to Davies in the case of ordinary quantum mechanics [23]. Although (*i*) classically the Equivalence Principle has a *local* character, and (*ii*) in a uniform gravitational field the Schrödinger equation leads to mass-dependent results, one can still say that the Equivalence Principle is realized at the quantum level if the motion of a quantum particle is considered to be described by a wave packet. This is a consequence of the Ehrenfest's theorem (see discussion in [23]). Nevertheless, Davies asks about the validity of the Equivalence Principle in the case of *delocalized* quantum states, such as *energy eigenstates*. These do not have classical counterparts associated to a (localized) particle with a well-defined trajectory, being of a different nature as compared to those localized states. In order to analyse this, Davies considered a variant of the (*gendanken*) experiment of Galileo at the leaning Tower of Pisa, with particles of different masses that would be vertically projected up in a uniform gravitational field. For classical particles, in the neighbourhood of the Earth, for instance, the out-and-back time is twice that spent in climbing, but for quantum particles there is a non-vanishing probability that they tunnel into the classically forbidden region. This special feature of the quantum theory might then give rise to a departure from the classical turnaround time, since a delay might ensue. In ordinary quantum mechanics, when one considers stationary, energy eigenstates, Davies [23] showed that, when the measurement is made far from the classical turning point, the time of flight of a quantum particle is identical to the classical result. In this sense the (weak) Equivalence Principle is preserved in quantum mechanics. In order to measure the time of flight associated with *delocalized* states, Davies applied a simple model of *quantum clock*, due to Peres [25] (see Sec. 6). This clock runs just when the particle travels within a given region of interest and does not measure absolute instants of time, but just the time difference. This is crucial, as in such a way one can avoid the collapse of the wave function. Peres showed that his quantum clock model correctly records the time of flight of a free particle.

It is in the sense mentioned above that in this paper we show that the Equivalence Principle can be extended to the NCQM in the case of stationary, energy eigenstates. Nevertheless, we remark that we achieve this result by explicitly solving the (stationary) scattering problem of a quantum particle in the presence of a uniform gravitational field. Our approach to the problem does not closely follow that of Davies. We closely follow the original approach by Peres [25] instead and explicitly consider that the quantum clock is coupled to the particle. Then, we will show that, in the case of small interaction between the particle and the clock, and when the measurement is made far from the turning point, the time of flight will be given by the *phase shift* of the wave function *and* this shift correctly codifies the time as measured by the quantum clock. On the other hand, Davies does not consider in the Hamiltonian of the problem the coupling term between the particle and the clock. He applies an expression from which the time of flight is given by the derivative of the phase change of the particle's wave function due to the action of the gravitational field only, without explicitly considering the clock. We will discuss more on the contrast of both approaches at the end of Sec. 7.

This paper is organized as follows: in Sec. 2, we review the algebraic approach to NCQM on a plane; in Sec. 3, we review the ordinary gravitational quantum well; in Sec. 4, we consider the non-commutative gravitational quantum well; in Sec. 5, we establish an upper-bound on the non-commutativity parameter; in Sec. 6, we review the Peres quantum clock model and in Sec. 7 we apply it to investigate the time of flight of particle subjected to a uniform gravitational field in NCQM.

2 Quantum mechanics on a non-commutative plane

Quantum mechanics inspired the idea of space-time non-commutativity, which can be introduced by (see, for instance, [3])

$$[\hat{x}_\mu, \hat{x}_\nu] = i\theta_{\mu\nu} \quad , \tag{1}$$

where $\hat{x}$ are the *coordinate operators* (the non-commutative analogues of the ordinary coordinate functions) and $\theta_{\mu\nu}$, the non-commutativity parameter, is assumed here to be a constant, anti-symmetric, real-valued $d \times d$ matrix (d is the space-time dimension), with dimension of length squared. Thus one can think of $\sqrt{\theta}$ as the caractheristic scale of length involved in the uncertainty on the measurement of coordinates. The commutation relations (1) lead to the space-time uncertainty relations $\Delta x_\mu \Delta x_\nu \geq |\theta_{\mu\nu}|/2$. It follows that non-commutativity replaces a space-time point by a cell whose dimension is given by the Planck area.

In this paper we are concerned about the effect of space-space non-commutativity in a two-dimensional, balistic experiment, so that we set $\theta_{0j} = \theta_{j0} = 0$ and restrict the discussion to the Moyal plane, $\mathcal{A}_\theta(\mathbb{R}^2)$, the non-commutative space generated by the non-commutative coordinates $\hat{x}_1$ and $\hat{x}_2$ [12], with commutation relations given by

$$[\hat{x}_i, \hat{x}_j] = i\theta_{ij} \quad , \tag{2}$$

where $i, j = 1, 2$. In two dimensions, $\theta_{ij} = \theta\epsilon_{ij}$, where ϵ_{ij} is the Levi-Civita tensor.

A generic element of $\mathcal{A}_\theta(\mathbb{R}^2)$ can be written as

$$\hat{\psi} = \int_{\mathbb{R}^2} d^2k \, \tilde{\psi}(\vec{k}) \, e^{i(k_1\hat{x}_1 + k_2\hat{x}_2)} \quad , \tag{3}$$

where $\tilde{\psi}(\vec{k})$ is some complex valued function satisfying the necessary conditions to guarantee the existence of $\hat{\psi}$.

The Moyal plane it is to be considered as the non-commutative analogue of the space of wave functions of ordinary quantum mechanics. According to this point of view, equation (3) is the analogue of the plane wave decomposition of wave functions.

2.1 Observables and dynamics

The non-commutative coordinates act on the Moyal plane as multiplication operators. Differently from the ordinary case, in the NCQM the multiplications from the left and right lead to different results. Extending the left-action by linearity we get an algebra representation of $\mathcal{A}_\theta(\mathbb{R}^2)$ on itself (the "left-representation")

$$\hat{\alpha}^L : \mathcal{A}_\theta(\mathbb{R}^2) \to \mathcal{A}_\theta(\mathbb{R}^2) \quad , \quad \hat{\alpha}^L \hat{\psi} = \hat{\alpha}\hat{\psi} \quad , \tag{4}$$

for all $\hat{\alpha}$, $\hat{\psi} \in \mathcal{A}_\theta(\mathbb{R}^2)$.

In this non-commutative framework, the momenta $\hat{P}_1$ and $\hat{P}_2$ canonically conjugated to the operators $\hat{x}_1^L$ and $\hat{x}_2^L$ are defined as

$$\hat{P}_1 = \frac{\hbar}{\theta}[\hat{x}_2, \cdot] \quad \text{and} \quad \hat{P}_2 = -\frac{\hbar}{\theta}[\hat{x}_1, \cdot] \quad , \tag{5}$$

the action on the Moyal plane generators being the expected one, *i.e.*,

$$\hat{P}_i \hat{x}_j = -i\hbar\delta_{ij} \quad . \tag{6}$$

One can show that the momenta $\hat{P}_1$ and $\hat{P}_2$ commute. Let $\hat{\psi}$ be a generic element of the Moyal plane. Then, by using the Jacoby identity (fulfilled by the commutator) we find

$$[\hat{P}_1, \hat{P}_2]\hat{\psi} = 0 \quad . \tag{7}$$

The commutation relations between $\hat{P}_i$ and $\hat{x}_j^L$ can be calculated directly. For example, we have

$$[\hat{P}_1, \hat{x}_1^L]\hat{\psi} = -i\hbar\hat{\psi} \quad . \tag{8}$$

Thus we can write the following generalized set of commutation relations

$$[\hat{x}_i^L, \hat{x}_j^L] = i\theta_{ij} \quad , \quad [\hat{P}_i, \hat{P}_j] = 0 \quad , \quad [\hat{P}_i, \hat{x}_j^L] = -i\hbar\delta_{ij} \quad . \tag{9}$$

We remark that our generalized commutation relations are distinct from the ones used in [19]-[22]. In [19]-[22] the momenta do not commute with the coordinates. And according to [19] such a phase-space non-commutativity gives rise to a modification of the Planck's constant, whereas [20] argues that this is not needed. Nevertheless, according to [24], those approaches are in fact physically equivalent, differing only in the manner one defines the non-commutative parameters.

Dynamics in non-commutative non-relativistic quantum mechanics is governed by the non-commutative Schrödinger equation

$$i\hbar \frac{\partial \hat{\psi}}{\partial t} = \hat{H} \hat{\psi} \quad . \tag{10}$$

In the simple cases the Hamiltonian operator $\hat{H}$ can be written in terms of the operators $\hat{x}_i^L$ and $\hat{P}_j$

$$\hat{H} = \hat{H}(\hat{x}_1^L, \hat{x}_2^L, \hat{P}_1, \hat{P}_2) \quad . \tag{11}$$

The action of $\hat{H}$ on wave functions belonging to the Moyal plane is obtained from (4) and (5).

2.2 Weyl-Moyal correspondence

In this section we introduce a useful correspondence between operators like in equation (3) and ordinary functions, the *Weyl-Moyal correspondence.*

The correspondence is the following: for every operator

$$\hat{\psi} = \int_{\mathbb{R}^2} d^2k \, \tilde{\psi}(\vec{k}) \, e^{i(k_1 \hat{x}_1 + k_2 \hat{x}_2)} \in \mathcal{A}_\theta(\mathbb{R}^2)$$

we associate a complex valued function of two variables, given by

$$\psi(x, y) := \int_{\mathbb{R}^2} d^2k \, \tilde{\psi}(\vec{k}) \, e^{i(k_1 x + k_2 y)} \quad , \tag{12}$$

which is called the *symbol* of $\hat{\psi}$. We denote this correspondence by $\mathcal{S}$, so that $\psi = \mathcal{S}(\hat{\psi})$.

Since the Moyal plane in an algebra, it is important to know the result of the Weyl-Moyal correspondence on a product of operators. Let $\hat{\psi}$ and $\hat{\phi}$ be two generic elements of $\mathcal{A}_\theta(\mathbb{R}^2)$. Using the Baker-Campbell-Hausdorff relation the product $\hat{\psi}\hat{\phi}$ can be written as

$$\hat{\psi}\hat{\phi} = \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} d^2k \, d^2p \, \tilde{\psi}(\vec{k}) \, \tilde{\phi}(\vec{p}) \, e^{-\frac{i\theta}{2}(k_1 p_2 - k_2 p_1)} \, e^{i[(k_1+p_1)\hat{x}_1 + (k_2+p_2)\hat{x}_2]} \quad . \tag{13}$$

Thus the symbol $\mathcal{S}(\hat{\psi}\hat{\phi})$ of $\hat{\psi}\hat{\phi}$ is the function

$$\mathcal{S}(\hat{\psi}\hat{\phi})(x,y) = \int_{\mathbb{R}^2}\int_{\mathbb{R}^2} d^2k\, d^2p\, \tilde{\psi}(\vec{k})\, \tilde{\phi}(\vec{p})\, e^{-\frac{i\theta}{2}(k_1p_2-k_2p_1)}\, e^{i[(k_1+p_1)x+(k_2+p_2)y]} \quad . \tag{14}$$

We note that the symbol $\mathcal{S}(\hat{\psi}\hat{\phi})$ can be equivalently written as

$$\mathcal{S}(\hat{\psi}\hat{\phi})(x,y) = \psi(x,y)\, \exp\left[\frac{i\theta}{2}\left(\frac{\overleftarrow{\partial}}{\partial x}\frac{\overrightarrow{\partial}}{\partial y} - \frac{\overleftarrow{\partial}}{\partial y}\frac{\overrightarrow{\partial}}{\partial x}\right)\right]\, \phi(x,y) \quad , \tag{15}$$

where $\psi(x,y)$ and $\phi(x,y)$ are the symbols of $\hat{\psi}$ and $\hat{\phi}$, respectively. In (15) the left (right) arrow means that one should differentiate the function on the left (right) side of the derivative symbol.

The Eq. (15) suggests the definition of an associative product between functions. It is called the Grönewold-Moyal product (in two dimensions),

$$(\psi \star \phi)(x,y) := \psi(x,y)\, \exp\left[\frac{i\theta}{2}\left(\frac{\overleftarrow{\partial}}{\partial x}\frac{\overrightarrow{\partial}}{\partial y} - \frac{\overleftarrow{\partial}}{\partial y}\frac{\overrightarrow{\partial}}{\partial x}\right)\right]\, \phi(x,y) \quad . \tag{16}$$

The associativity of this $\star$-*product* follows from the associativity of the product between elements of the Moyal plane.

One can verify by direct calculation that the Weyl-Moyal correspondence is compatible with the momenta, *i.e.*,

$$\mathcal{S}(\hat{P}_1\hat{\psi}) = -i\hbar\frac{\partial\psi}{\partial x} \quad \text{and} \quad \mathcal{S}(\hat{P}_2\hat{\psi}) = -i\hbar\frac{\partial\psi}{\partial y} \quad , \tag{17}$$

so that the kinetic part of the Schrödinger equation is preserved under $\mathcal{S}$. In order to write the image of the Schrödinger equation under the action of $\mathcal{S}$ we still have to find out how the potential energy term will look like after we apply the correspondence $\mathcal{S}$ on it. For this purpose we make use the definition of the Grönewold-Moyal product to write $\mathcal{S}(\hat{V}\hat{\psi}) = V \star \psi$. Thus we have

$$-\frac{\hbar^2}{2m}\bigtriangledown^2\psi + V\star\psi = i\hbar\frac{\partial\psi}{\partial t} \quad , \tag{18}$$

since the time derivative commutes with $\mathcal{S}$.

Now, by seting $\psi(\vec{r},t) = \psi(\vec{r})\, e^{-\frac{i}{\hbar}Et}$, the time-independent equation is

$$-\frac{\hbar^2}{2m}\bigtriangledown^2\psi + V\star\psi = E\psi \quad , \tag{19}$$

where E is the energy of the particle and ψ is time-independent.

3 The gravitational quantum well

In this section we review the problem of a quantum neutral particle of inertial mass m travelling in the ordinary (commutative) xy plane. The particle is under the influence of a uniform gravitational field in the vertical direction (Oy), while in the horizontal direction (Ox) it is free. Besides that, we consider a perfectly reflecting mirror at the bottom ($y = 0$), so that the particle gets trapped. We will summarise the main facts concerning the energy spectrum of this system and set up the notation to be used in future sections.

Let $\vec{g}$ be a uniform gravitational field and let the Oy axis be oriented in the opposite direction of $\vec{g}$, *i.e.*, $\vec{g} = -g\,\hat{e}_y$. The potential energy associated to the particle is $V(x,y) = mgy$. At the boundary ($y = 0$) the particle encounters non-gravitational forces, which prevent it to get into the negative portion of the Oy axis. We can thus write

$$V(x,y) = \begin{cases} mgy\,, & y > 0\,, \quad \forall x \quad , \\ \infty\,, & y = 0\,, \quad \forall x \quad . \end{cases} \tag{20}$$

The time-independent Schrödinger equation for a non-relativistic particle under the action of $V(x,y)$ reads

$$-\frac{\hbar^2}{2m}\bigtriangledown^2\psi(x,y) + V(x,y)\psi(x,y) = E\psi(x,y) \quad , \tag{21}$$

where E is the total energy of the particle.

The solution of the energy eigenvalue problem implied by (21), the so called *gravitational quantum well*, is well-known and treated in some textbooks and pedagogic articles (for instance, see [15]).

Since the particle is trapped along the vertical direction, its energy spectrum has a discrete part associated to this degree of freedom. On the other hand, its free motion in the horizontal direction gives rise to a continuous contribution to its energy spectrum. The part of the energy spectrum related to the vertical direction gives the gravitational quantum bound states of the particle. Let us now briefly review how this comes about.

The free particle potential in the horizontal direction suggests we use the *ansatz*

$$\psi(x,y) = e^{ikx}\phi(y) \tag{22}$$

in the time-independent Schrödinger equation, which implies that the function ϕ satisfies the Airy equation [26]

$$\frac{d^2\phi(\xi)}{d\xi^2} - \xi\phi(\xi) = 0 \quad , \tag{23}$$

where $\xi = (\frac{y-b}{a})$, $a = \left(\frac{\hbar^2}{2m^2 g}\right)^{1/3}$ and $b = \frac{E - \frac{\hbar^2 k^2}{2m}}{mg}$.

The two linear independent solutions of (23) are denoted by $Ai(\xi)$ and $Bi(\xi)$. The behaviour of these functions is shown in Fig. 1.



Figure 1: *The Airy functions $Ai(\xi)$ (bounded as $\xi \to \infty$) and $Bi(\xi)$ (unbounded as $\xi \to \infty$). The classical turning point corresponds to $\xi = 0$.*

The general solution of (23) then reads

$$\phi(y) = A\,Ai\left(\frac{y-b}{a}\right) + B\,Bi\left(\frac{y-b}{a}\right) \quad , \tag{24}$$

where A, B are coefficients to be determined by boundary conditions. The restriction imposed by the perfectly reflecting mirror at the bottom implies that $\psi(x,y) = 0$ for $y = 0$. Besides that, the wave function $\phi(y)$ corresponding to the vertical motion is a bound state, which means that the probability of finding the particle far from the bottom is vanishingly small. The boundary conditions relevant to the problem are thus the following

$$\begin{array}{lll} \phi \to 0 \quad , & y \to \infty \quad , \\ \phi = 0 \quad , & y = 0 \quad . \end{array} \tag{25}$$

The first one gets rid of $Bi(\xi)$, since it diverges as $y \to \infty$. The second boundary condition leads to a quantization condition for the energy eigenvalues associated to ϕ, so that they can be written in terms of the zeroes of the Airy function. The final form of the particle energy spectrum reads

$$E_{k,n} = \frac{\hbar^2 k^2}{2m} + \left(\frac{mg^2\hbar^2}{2}\right)^{1/3} . (-\alpha_n) \quad , \tag{26}$$

where α_n is the n-th zero of the Airy function Ai [26].

4 The non-commutative gravitational quantum well

In this section we consider the non-commutative version of the problem discussed earlier. The calculations go easier if we work directly with functions, *i.e.*, if we apply the Weyl-Moyal correspondence in both sides of the Schrödinger equation, just like we did in Sec. 2.2.

By applying the product (16) to the case of the gravitational potential $V(x,y) = mgy$ we find

$$V(x,y) \star \psi(x,y) = mgy \star \psi(x,y) = mg\left(y\psi(x,y) - \frac{i\theta}{2}\frac{\partial\psi}{\partial x}\right) \quad , \tag{27}$$

so that the Schrödinger equation reads

$$-\frac{\hbar^2}{2m}\frac{\partial^2\psi(x,y)}{\partial x^2} - \frac{\hbar^2}{2m}\frac{\partial^2\psi(x,y)}{\partial y^2} + mgy\psi(x,y) - \frac{mgi\theta}{2}\frac{\partial\psi}{\partial x} = E\psi(x,y) \quad . \tag{28}$$

The simplicity of the result (27) comes from the fact that the gravitational potential is a linear function of y only, so that the Grönewold-Moyal product series (Eq. (16)) finishes at its first order term.

As the motion is free along Ox, we use the *ansatz* $\psi(x,y) = e^{ikx}\phi(y)$ into (28) to get

$$\frac{d^2\phi}{dy^2} + \frac{2m^2 g}{\hbar^2}\left(\frac{\mathcal{E}_\theta}{mg} - y\right)\phi(y) = 0 \quad , \tag{29}$$

where

$$\mathcal{E}_\theta = E - \frac{\hbar^2 k^2}{2m} - \frac{mgk\theta}{2} \quad . \tag{30}$$

Now, by setting

$$\xi = \frac{y - b_\theta}{a} \quad , \quad a = \left(\frac{\hbar^2}{2m^2 g}\right)^{1/3} \quad \text{and} \quad b_\theta = \frac{\mathcal{E}_\theta}{mg} \quad , \tag{31}$$

we can put (29) in the form of an Airy equation

$$\frac{d^2\phi(\xi)}{d\xi^2} - \xi\phi(\xi) = 0 \quad , \tag{32}$$

whose general solution is

$$\phi(y) = A\, Ai\left(\frac{y - b_\theta}{a}\right) + B\, Bi\left(\frac{y - b_\theta}{a}\right) \quad . \tag{33}$$

The boundary conditions for $\phi(y)$ are of the same type as in (25), *i.e.*,

$$\phi = 0 \quad , \quad y = 0 \quad , \tag{34}$$

$$\phi \to 0 \quad , \quad y \to \infty \quad . \tag{35}$$

From condition (35) we are left with

$$\phi(y) = A\, Ai\left(\frac{y - b_\theta}{a}\right) \quad . \tag{36}$$

Now, from the boundary condition (34), it follows that $Ai\left(-\frac{b_\theta}{a}\right) = 0$, so that $-b_\theta/a$ are the roots of the Airy function Ai, *i.e.*,

$$b_{\theta,n} = -a\,\alpha_n \quad , \tag{37}$$

where α_n denotes the n-th zero of Ai. The result (37) combined with the definition of b_θ (see Eq.'s (30) and (31)) gives the spectrum of the Hamiltonian

$$E_{k,n,\theta} = \frac{\hbar^2 k^2}{2m} + \left(\frac{mg^2\hbar^2}{2}\right)^{1/3} .\, (-\alpha_n) + \frac{mgk\theta}{2} \quad . \tag{38}$$

For $\theta = 0$, the result of ordinary quantum mechanics (Eq. (26)) is recovered. We note that the effect of non-commutativity also disappears for $k = 0$, what corresponds simply to have a one-dimensional movement, along the direction of the gravitational field. Hence, the above spectrum reflects the non-commutative nature of the configuration space.

We see that the spectrum is not invariant by changing k into $-k$. In fact, it is easy to show that the parity transformation does not leave the Hamiltonian spectrum invariant (one should recall that in two spatial dimensions the change $x \longrightarrow -x$, while y is kept fixed, is a parity transformation). This is in accordance with the fact that the parity transformation is not a symmetry of the commutation relation between the coordinate operators.

The Eq. (38) is the main result of this section. In the next section we connect this formula with the available experimental data of the Grenoble experiment. This will allow us to impose an upper-bound on the value of the non-commutative constant θ.

We finish this section by analysing what is, now in NCQM, the analogue of the classical turning point in ordinary quantum mechanics. From Eq. (31) we see that $\xi = 0$ implies $y = b_\theta$. In order to understand the meaning of this, let us first consider the case when $\theta = 0$, for which $\xi = 0$ implies $y = b = \frac{E - \frac{\hbar^2 k^2}{2m}}{mg}$, the classical turning point. Now, for $\theta \neq 0$, the point $b_\theta = \frac{\mathcal{E}_\theta}{mg}$ does not correspond to the turning point of the classical motion. In spite of this, it still corresponds to the point from where the wave function exponentially decays. This suggests that in NCQM the forbidden region has a distinct nature than that in ordinary quantum mechanics. A through analysis of the probability density in NCQM in general will be helpful in completely clarifying that point. Efforts in this direction are under development.

5 The Grenoble experiment and an upper-bound on θ

Having found the bound states of a particle in a non-commutative gravitational quantum well, we can now constraint the value of the non-commutative parameter, by resorting to data of the experiment performed by Nesvizhevsky *et al.* - the *Grenoble experiment* [16]-[18]. In this experiment the lowest quantum bound states of neutrons on free-fall in the Earth's gravitational field were observed and their energy determined. Due to the weaker strength of gravity, as compared to nuclear and electromagnetic interactions, this experiment provided another manifestation of the quantum nature of matter, but now under challeging, extremely sensible experimental designed conditions, being another striking observation of the wave behaviour of matter in a gravitational field [1]. This was achieved by means of a gravitational quantum well, formed by the Earth's gravitational field and a horizontal reflecting mirror (considered as perfectly reflecting). Due to their charge neutrality, long lifetime, and low mass, neutrons are suitable to perform this kind of experiment, in which the effect of the interactions other than gravity must be negligible. A horizontal beam of ultracold neutrons was thus allowed to fall freely, flying above the reflecting mirror at the bottom. As no forces act on neutrons horizontally and just gravity acts vertically, we have a gravitational potential well along the latter direction. For details on the experimental setup, see [16]-[18].

An upper-bound on θ can be established by imposing that the θ-dependent corrections to the energy implied by Eq. (38) be smaller or of the order of the maximum differences of the energy levels provided by the Grenoble experiment, *i.e.*, according to its error bars. Hence we require that

$$\frac{mgk\theta}{2} \lesssim \Delta E^{exp} \quad , \tag{39}$$

so that

$$\theta \lesssim \frac{2\Delta E^{exp}}{mgk} \quad . \tag{40}$$

For the first two energy levels, the experiment gives $\Delta E_1^{exp} = 6.55 \times 10^{-32}\, J \quad (n = 1)$ and $\Delta E_2^{exp} = 8.68 \times 10^{-32}\, J \quad (n = 2)$. For neutrons we have $m \simeq 1.675 \times 10^{-27}\, kg$, and in the experiment the neutrons had a mean horizontal speed $< v_x > \simeq 6.5\, m/s$ [18], so that $k = \frac{<p_x>}{\hbar} = \frac{m<v_x>}{\hbar} \simeq 1.03 \times 10^8\, m^{-1}$. Then by considering $g \simeq 9.81\, m/s^2$ we arrive at the following upper-bounds for θ

$$\begin{aligned} \theta &\lesssim 0.771 \times 10^{-13}\, m^2 \qquad (n = 1) \quad , & (41)\\ \theta &\lesssim 1.021 \times 10^{-13}\, m^2 \qquad (n = 2) \quad . & (42) \end{aligned}$$

As far as we know, other works which resort to data of the Grenoble experiment do not arrive at un upper-bound on θ. As they considered non-commutativity in both configuration and momentum spaces, they could find an upper-bound on the parameter of momentum non-commutativity instead. [19]-[22]. Since we consider spatial non-commutativity only, we can establish an upper-bound on the parameter of spatial non-commutativity. We remark that other upper-bounds on θ were established by experimental data related to the Lorentz invariance [28] and to the lamb shift [29], for instance.

---

[1] An earlier experiment in which the quantum, wave nature of matter was manifested, due to its interaction with the Earth's gravitational field, is that of the observation of gravitationally induced quantum interference of neutrons [27].

One of the most important prospects about the Grenoble experiment is the improvement of the energy resolution of the neutrons energy levels. In principle, with the energy resolution, ΔE, limited only by the uncertainty principle ($\Delta E\,\Delta t \ \sim \hbar$), one could achieve a value as low as $10^{-18} eV$, if Δt approaches the neutron lifetime. Improvement in the energy resolution could help in testing the proportionality between inertial and gravitational masses for neutrons [16]-[18]. The Equivalence Principle in the context of the gravitational quantum well was studied in [30], but in a manner completely different from that we consider in the following section. In fact, in the following, we do not even need to consider that a well is set up.

6 Quantum clock and time of flight measurement

By resorting to the basic notion that "a clock is a dynamical system which pass through a succession of states at constant time intervals" [25], that is, that "... the measurement of time actually is the observation of some dynamical variable, the law of motion of which is known ..." [25], Peres has modeled a simple quantum clock by means of a quantum rotor (also known as the Larmor clock). When coupled to another system, this can measure the duration of physical processes as well as keep a permanent record of it, such as the time of flight of a particle, or even to control the duration of a physical process [25] (see also [31]).

In order to address the essential features involved in modeling a quantum clock, in the following we briefly review Peres construction of a quantum clock and its particular use in a time of flight experiment involving the motion of a free quantum particle, as considered in [25]. The features involved in this case will guide us in applying the Peres quantum clock to the determination of the time of flight of a quantum particle in a uniform gravitational field in the context of NCQM.

Peres basic ideia in modeling a quantum clock is to consider a quantum rotor, since this can be regarded as describing the motion of a pointer on a clock dial. Thus, he considered a clock with an odd number of "pointer states", $N = 2j + 1$, represented by

$$u_n(q) = \frac{1}{\sqrt{2\pi}} e^{inq} \quad , \qquad n = -j, ..., j \quad , \tag{43}$$

where q is the clock's degree of freedom and $0 \leq q < 2\pi$.

Another suitable orthogonal basis of states for the clock is

$$v_\kappa = \frac{1}{\sqrt{N}} \sum_{n=-j}^{j} e^{-i2\pi\kappa n/N} \, u_n(q) \quad , \tag{44}$$

$$v_\kappa = \frac{1}{\sqrt{2\pi N}} \frac{\sin \frac{N}{2}(q - \frac{2\pi\kappa}{N})}{\sin \frac{1}{2}(q - \frac{2\pi\kappa}{N})} \quad , \tag{45}$$

where $\kappa = 0, ..., N-1$. It follows that for large values of N the state $v_\kappa(q)$ is sharply picked at $\frac{2\pi\kappa}{N}$. We can refer to the states v_κ's as the clock states.

By defining the projection operators $P_\kappa v_n = \delta_{\kappa n} v_n$, a "clock time" operator can be defined as

$$T_c = \tau \sum_\kappa \kappa P_\kappa \quad , \tag{46}$$

where τ is the time resolution of the clock. Its eigenstates are v_κ, with $t_\kappa = \kappa\tau$ as the corresponding eigenvalues.

The clock's Hamiltonian is

$$H_c = \omega J \quad , \tag{47}$$

where $\omega = \frac{2\pi}{N\tau}$ and $J = -i\hbar\partial/\partial q$. The eigenstates of H_c are the vectors $u_n(q)$, defined in Eq. (43). The eigenvalue corresponding to $u_n(q)$ is $n\hbar\omega$.

From (43) and (47) it follows that

$$e^{-iH_c\tau/\hbar} v_\kappa(q) = v_{\kappa+1(mod\, N)}(q) \quad . \tag{48}$$

Assuming that v_0 is the initial state of the clock, the above result implies that the clock will pass successively through the states $v_0, v_1, v_2, ...$, at time intervals given by τ.

It is interesting to see the Peres construction applied in the simplest case before we go into the details of the gravitational case. Thus we next apply the Peres quantum clock to the determination of the time of flight of a free particle traveling between two assigned points [25]. One has to demand that the clock be activated when the particle pass by the first point and then be stopped when it passes by the second one. Let us consider that the particle moves along the direction Ox, so that x_1 and x_2 correspond to the two positions of interest. In the Schrödinger representation the Hamiltonian for the composed, particle-clock system reads

$$H = -\frac{\hbar^2}{2m}\frac{\partial^2}{\partial x^2} - i\hbar\omega\Theta(x-x_1)\Theta(x_2-x)\frac{\partial}{\partial q} \quad , \tag{49}$$

so that the clock runs only if the particle lies within the interval $[x_1, x_2]$.

Following Peres, we set the initial state of the clock as $v_0 = \sum_n \frac{u_n}{\sqrt{N}}$. As H and $H_c = \omega J$ commute, it is simpler to solve the equation of motion for the clock in an eingenstate of J, *i.e.*, u_n, and then sum over these partial solutions to get the evolved wave function corresponding to the initial state, v_0. Therefore, the initial state of the (still not coupled) particle-clock system can be written as $\Psi_0(x,q)e^{-iEt/\hbar}$, with $\Psi_0(x,q)$ given in the factorized form (see [25, 32])

$$\Psi_0(x,q) = A_k\, e^{ikx} v_0(q) = A_k\, e^{ikx} \frac{1}{\sqrt{N}} \sum_{n=-j}^{j} u_n(q) \quad , \tag{50}$$

where E is the energy of the particle, $k = \sqrt{2mE}/\hbar$ and A_k is a normalization constant.

On the other hand, the final state of the system cannot be factorized as $\Psi_0(x,q)$, since the motion of the particle in the region $x_1 < x < x_2$ will activate the clock, making the particle and clock coordinates to mix. Thus we write

$$\Psi(x,q) = \frac{1}{\sqrt{N}} \sum_{n=-j}^{j} \psi_k^n(x) u_n(q) \quad . \tag{51}$$

The substitution of (51) into the Schrödinger equation leads to

$$-\frac{\hbar^2}{2m}\frac{d^2\psi_k^n}{dx^2} + \Big[n\hbar\omega\Theta(x-x_1)\Theta(x_2-x) - E\Big]\psi_k^n(x) = 0 \quad . \tag{52}$$

Thus we see that ψ_k^n satisfies the Schrödinger equation for a particle under the action of a rectangular barrier of height $V_n = n\hbar\omega$ and length

$L = x_2 - x_1$. Outside the barrier we have (we are basically following Leavens presentation of the original Peres quantum clock model [32])

$$\psi_k^n(x) = \begin{cases} e^{ikx} + R_n(k)\, e^{-ikx} & , \quad x < x_1 \quad , \\ T_n(k)\, e^{ikx} & , \quad x > x_2 \quad , \end{cases} \tag{53}$$

where $T_n(k)$ and $R_n(k)$ are the transmission and reflexion amplitudes, respectively, given by

$$T_n(k) = \frac{e^{-ikL}}{\cos(k_n L) - \frac{i}{2}\left(\frac{k}{k_n} + \frac{k_n}{k}\right)\sin(k_n L)} \tag{54}$$

and

$$R_n(k) = \frac{\frac{i}{2}\left(\frac{k_n}{k} - \frac{k}{k_n}\right) e^{2ikx_1}\sin(k_n L)}{\cos(k_n L) - \frac{i}{2}\left(\frac{k}{k_n} + \frac{k_n}{k}\right)\sin(k_n L)} \quad , \tag{55}$$

and where $k_n = \frac{\sqrt{2m(E - n\hbar\omega)}}{\hbar}$.

Of course that the particle will be perturbed when coupled to the physical clock. In order that one has a small disturbance on particle's evolution, one must consider that such coupling is sufficiently small. We therefore set $V_n \ll E$, from which we have $k_n \simeq k - n\omega(2E/m)^{-1/2} = k - n\omega/(\hbar k/m)$. Besides that, in the limit of small coupling we also have

$$T_n(k) \simeq \exp\left(i(k_n - k)L\right) \quad , \tag{56}$$

so that $|T_n(k)| \simeq 1$. Under the above conditions we can write the phase shift due to the clock as

$$(k_n - k)L \simeq -n\omega L(2E/m)^{-1/2} = -\frac{n\omega L}{\hbar k/m} \quad . \tag{57}$$

Let us call $T = L/(\hbar k/m)$. Now we can express the final wave function for the particle-clock system as

$$\Psi(x,q) \simeq A_k e^{ikx} \frac{1}{\sqrt{N}} \sum_{n=-j}^{j} e^{-in\omega L/(\hbar k/m)} u_n(q) \quad , \tag{58}$$

i.e.,

$$\Psi(x,q) \simeq A_k e^{ikx} \frac{1}{\sqrt{N}} \sum_{n=-j}^{j} e^{-in\omega T} u_n(q) = A_k e^{ikx} v_0(q - \omega T) \quad . \tag{59}$$

Since $e^{-i\hat{H}_c T/\hbar} u_n(q) = u_n(q - \omega T)$, it follows that the pointer, initially directed to $q = 0$, will be found directed to $q \simeq \omega T$ after the particle leaves the region where the clock runs. By realizing that $v = \sqrt{2E/m} = \hbar k/m$ is the velocity of a free quantum particle with energy E, we see that the Peres clock records the time of flight of a particle which travels between the two assigned points. The key fact is that the time of flight is encoded in the phase shift due to the clock barrier.

We remark that the quantum clock does not measure the absolute instants of time in which the particle passes through the positions x_1 and x_2, but only the time difference to travel between them. This fact avoids the collapse of the particle wave function. That is the advantage of using a quantum clock to study delocalized states, as energy eigenstates (see Sec. 7).

7 Particle in a uniform gravitational field: non-commutative case

In [23] Davies applied the Peres model of a quantum clock to determine the time of flight of a quantum particle in its round trip in a uniform gravitational field, when it is vertically projected up. Davies interest on studying this sort of "quantum Galileo experiment" was motivated by asking if there would be a violation of the (Weak) Equivalence Principle when one considers quantum particles, as they may tunnel into the classically forbidden region of the gravitational potential. Therefore, a mass-dependent delay in the time of flight might result, as compared to the classical case. As Davies remarks, the answer to this question is not *a priori* obvious in the case one is handling with energy eigenstates, which are delocalized states, with no corresponding classical counterparts on localized bodies (for a study of the motion of wave packets and the "quantum Galileo experiment", see [33]). Davies finds, by assuming the equality of the inertial (m_i) and gravitational (m_g) masses, that when the measurement is made far from the classical turning point, the time of flight is equal to the classical result. Therefore, in this sense, the (Weak) Equivalence Principle holds in quantum mechanics, even when one considers highly non-classical states.

Inspired by Davies, we next apply the quantum clock model of Peres to determine the time of flight of a particle in a uniform gravitational field in the context of NCQM, in order to address the "quantum Galileo experiment" and the question of the validity of the Equivalence Principle. We will assume that $m_i = m_g \equiv m$.

We make use of the simple clock model of Sec. 6 to measure the interval of time during which the particle lies within the semi-infinite region of space defined by $y_0 \leq y < \infty$, where y_0 is some fixed height, from where we can consider that the clock is set up. We consider that the particle is obliquely projected up from some height below y_0.

Let P_{y_0} be the projection operator defined by

$$P_{y_0}\psi(x,y) = \Theta(y-y_0)\psi(x,y) \quad . \tag{60}$$

The vertical particle's degree of freedom couples to that of the quantum clock through the interaction Hamiltonian

$$H_I = P_{y_0}H_c = \Theta(y-y_0)\omega J \quad . \tag{61}$$

Of course that H_I is a tensor product of operators, since P_{y_0} and H_c act on two independent Hilbert spaces.

We now consider the operator $\hat{P}_{y_0} = \Theta(\hat{x}_2 - y_0)$, the non-commutative analogue of P_{y_0}. We know from Sec. 2.2 that the action of the potential energy $\hat{V}(\hat{x}_1, \hat{x}_2)$ on non-commutative wave functions is mapped into a Moyal product under the Weyl-Moyal correspondence. Denoting the energy by E, the eigenstate of the particle-clock system is given by $\Psi(x,y,q)e^{-iEt/\hbar}$, so that the time-independent non-commutative Schrödinger equation reads

$$-\frac{\hbar^2}{2m}\frac{\partial^2\Psi}{\partial x^2} - \frac{\hbar^2}{2m}\frac{\partial^2\Psi}{\partial y^2} + mgy\star\Psi + \Theta(y-y_0)\star\omega J\Psi = E\Psi \quad . \tag{62}$$

The Grönewold-Moyal product of $V = mgy$ by $\Psi(x,y,q)$ is given by

$$mgy\star\Psi(x,y,q) = mg\left(y\Psi(x,y,q) - \frac{i\theta}{2}\frac{\partial\Psi(x,y,q)}{\partial x}\right) \quad , \tag{63}$$

just like in Eq. (27).

Substituting (63) into (62) we find

$$-\frac{\hbar^2}{2m}\frac{\partial^2\Psi}{\partial x^2} - \frac{\hbar^2}{2m}\frac{\partial^2\Psi}{\partial y^2} + mg\left(y\Psi - \frac{i\theta}{2}\frac{\partial\Psi}{\partial x}\right) + \Theta(y-y_0)\star\omega J\Psi = E\Psi \quad . \tag{64}$$

We now make use of the *ansatz* $\Psi(x,y,q) = e^{ikx}\,\Phi(y,q)$ into (64) to write

$$-\frac{\hbar^2}{2m}\frac{\partial^2\Psi}{\partial y^2} + mgy\,\Psi + \Theta(y-y_0)\star\omega J\Psi = \left(E - \frac{\hbar^2k^2}{2m} - \frac{mgk\theta}{2}\right)\Psi \quad . \tag{65}$$

It remains to calculate the product $\Theta(y-y_0)\star\omega J\Psi$. At first, notice that $\Theta(y-y_0)$ is a distribution. If we try to deal with it as we would do with an ordinary function, we have to face the fact that it is not differentiable, thus leading to the question of how to apply the definition (16) of the Grönewold-Moyal product to this case. It is reasonable to try just to extend the result that would be obtained in the case of a smooth function $F(y)$ to the case of $\Theta(y-y_0)$. By directly applying (16) we find

$$F(y)\star\Psi(x,y,q)=\sum_{n=0}^{\infty}\frac{1}{n!}\left(\frac{-i\theta}{2}\right)^{n}\frac{d^nF(y)}{dy^n}\frac{\partial^n\Psi(x,y,q)}{\partial x^n}\quad , \tag{66}$$

but since the x dependency of Ψ is just e^{ikx}, we get

$$F(y)\star\Psi=\left[\sum_{n=0}^{\infty}\frac{1}{n!}\left(\frac{k\,\theta}{2}\right)^{n}\frac{d^nF(y)}{dy^n}\right]\Psi=F\left(y+\frac{k\,\theta}{2}\right)\Psi\quad , \tag{67}$$

which lead us to write the following expression for $\Theta(y-y_0)\star\Psi(x,y)$

$$\Theta(y-y_0)\star\Psi(x,y,q)=\Theta(y-y_\theta)\Psi(x,y,q)\quad , \tag{68}$$

where

$$y_\theta=y_0-\frac{k\,\theta}{2}\quad . \tag{69}$$

In order to verify that (68) is right, one has to apply (in the distributional sense) both sides of (68) onto a test function $f(y)$ and check that

$$\int_{-\infty}^{\infty}dy\,\Theta(y-y_0)\star\Psi(x,y,q)\,f(y)=\int_{-\infty}^{\infty}dy\,\Theta(y-y_\theta)\Psi(x,y,q)\,f(y)\quad . \tag{70}$$

It turns out that (70) is correct, so that the above naive procedure is completely justified.

Substituting (68) into (65) and simplyfing the common factor e^{ikx} we get

$$-\frac{\hbar^2}{2m}\frac{\partial^2\Phi}{\partial y^2}+mgy\Phi+\Theta(y-y_\theta)\omega J\,\Phi=\left(E-\frac{\hbar^2k^2}{2m}-\frac{mgk\theta}{2}\right)\Phi\quad . \tag{71}$$

It is now useful to write the wave function $\Phi(y,q)$ just like in (51)

$$\Phi(y,q)=\frac{1}{\sqrt{N}}\sum_{n=-j}^{j}\psi_k^n(y)u_n(q)\quad , \tag{72}$$

where $u_n(q)$ are the basis vectors defined in (43), *i.e.*, $u_n(q) = \frac{1}{\sqrt{2\pi}}\, e^{inq}$.

Substituting (72) into (71) we find

$$\frac{1}{\sqrt{N}} \sum_{n=-j}^{j} \left\{ -\frac{\hbar^2}{2m} \frac{d^2\psi_k^n}{dy^2} + \left[mgy + n\hbar\omega\Theta(y-y_\theta) - \mathcal{E}_\theta \right] \psi_k^n(y) \right\} u_n(q) = 0 \quad , \tag{73}$$

where

$$\mathcal{E}_\theta = E - \frac{\hbar^2 k^2}{2m} - \frac{mgk\theta}{2} \quad . \tag{74}$$

But since the u_n are linearly independent, we can write a differential equation for each $\psi_k^n(y)$

$$-\frac{\hbar^2}{2m} \frac{d^2\psi_k^n}{dy^2} + \left[mgy + n\hbar\omega\Theta(y-y_\theta) - \mathcal{E}_\theta \right] \psi_k^n(y) = 0 \quad . \tag{75}$$

There are two different regions to consider, according to the value assumed by the Θ distribution. For $y < y_\theta$ we have $\Theta(y-y_\theta) = 0$, so that the Schrödinger equation reduces to

$$\frac{d^2\psi_k^n(y)}{dy^2} + \frac{2m^2 g}{\hbar^2} \left(\frac{\mathcal{E}_\theta}{mg} - y \right) \psi_k^n(y) = 0 \quad , \tag{76}$$

whose general solution (see Eq.'s (29)-(33)) is

$$\psi_k^n(y) = A\, Ai\left(\frac{y-b_\theta}{a} \right) + B\, Bi\left(\frac{y-b_\theta}{a} \right) \quad , \tag{77}$$

with

$$a = \left(\frac{\hbar^2}{2m^2 g} \right)^{1/3} \quad \text{and} \quad b_\theta = \frac{\mathcal{E}_\theta}{mg} = b - \frac{k\theta}{2} \quad . \tag{78}$$

For $y \geq y_\theta$ we have $\Theta(y-y_\theta) = 1$, so that the Schrödinger equation reads

$$\frac{d^2\psi_k^n(y)}{dy^2} + \frac{2m^2 g}{\hbar^2} \left(\frac{\mathcal{E}_\theta^n}{mg} - y \right) \psi_k^n(y) = 0 \quad , \tag{79}$$

where

$$\mathcal{E}_\theta^n = \mathcal{E}_\theta - n\hbar\omega \quad . \tag{80}$$

The Eq. (79) is analogous to Eq. (76), the only difference being in the value of the energy. It follows that its general solution is a linear combination of Airy functions too. Now, notice that the Airy function Bi is physically unacceptable in the positive semi-axis, since it diverges for large positive values of its argument. Taking this into account we write the general solution of the Schrödinger equation for $y \geq y_\theta$

$$\psi_k^n(y) = C\, Ai\left(\frac{y - b_\theta^n}{a}\right) \quad , \tag{81}$$

where

$$b_\theta^n = \frac{\mathcal{E}_\theta^n}{mg} = b_\theta - \frac{n\hbar\omega}{mg} \quad . \tag{82}$$

The complex constants in (77) and (81) are determined by the matching conditions at $y = y_\theta$. By setting $\xi = \frac{y - b_\theta}{a}$, we see that $y = y_\theta \longrightarrow \xi = \frac{y_\theta - b_\theta}{a} = \frac{y_0 - b}{a}$. In what follows we denote this particular value of ξ by the symbol ξ_0. Thus,

$$A\, Ai\left(\xi_0\right) + \, B\, Bi\left(\xi_0\right) = C\, Ai\left(\xi_0 + \frac{n\hbar\omega}{mga}\right) \quad , \tag{83}$$

$$A\, \frac{dAi}{d\xi}\left(\xi_0\right) + \, B\, \frac{dBi}{d\xi}\left(\xi_0\right) = C\, \frac{dAi}{d\xi}\left(\xi_0 + \frac{n\hbar\omega}{mga}\right) \quad . \tag{84}$$

By solving this system we find the values of $\frac{B}{A}$ and $\frac{C}{A}$, namely

$$\frac{B}{A} = \frac{C}{A}\, \frac{Ai\left(\xi_0 + \frac{n\hbar\omega}{mga}\right)}{Bi\left(\xi_0\right)} - \frac{Ai\left(\xi_0\right)}{Bi\left(\xi_0\right)} \quad , \tag{85}$$

$$\frac{C}{A} = \frac{Ai\left(\xi_0\right)\, \frac{dBi}{d\xi}\left(\xi_0\right) - \frac{dAi}{d\xi}\left(\xi_0\right)\, Bi\left(\xi_0\right)}{Ai\left(\xi_0 + \frac{n\hbar\omega}{mga}\right)\, \frac{dBi}{d\xi}\left(\xi_0\right) - \frac{dAi}{d\xi}\left(\xi_0 + \frac{n\hbar\omega}{mga}\right)\, Bi\left(\xi_0\right)} \quad . \tag{86}$$

The constant A remains undetermined. We will choose its value later.

Notice that the numerator in (86) is the Wronskian of $Ai(\xi)$ and $Bi(\xi)$ at $\xi = \xi_0$. By using the fact that (see [26])

$$W[Ai(\xi), Bi(\xi)] = \frac{1}{\pi} \quad , \qquad \forall\, \xi \, , \tag{87}$$

we can write

$$\frac{C}{A} = \frac{1}{\pi}\, \frac{1}{Ai\left(\xi_0 + \frac{n\hbar\omega}{mga}\right)\, \frac{dBi}{d\xi}\left(\xi_0\right) - \frac{dAi}{d\xi}\left(\xi_0 + \frac{n\hbar\omega}{mga}\right)\, Bi\left(\xi_0\right)} \quad . \tag{88}$$

At this point it is useful to put together the results just obtained and write the complete solution of (62). At first we recall that $\Psi(x, y, q) = e^{ikx}\Phi(y, q)$. Besides that, Φ was decomposed according to (72). We thus write

$$\Psi_{E,k}(x, y, q) = \mathcal{C}_{E,k}\, \frac{e^{ikx}}{\sqrt{N}} \sum_{n=-j}^{j} \psi_k^n(y)\, u_n(q) \quad , \tag{89}$$

where

$$\psi_k^n(y) = \begin{cases} A\, Ai\, \left(\frac{y-b_\theta}{a}\right) + B\, Bi\left(\frac{y-b_\theta}{a}\right) & , \text{if} \quad y \leq y_\theta \\ C\, Ai\left(\frac{y-b_\theta^n}{a}\right) & , \text{if} \quad y \geq y_\theta \end{cases} \tag{90}$$

with B/A and C/A given by (85) and (88). $\mathcal{C}_{E,k}$ is a normalization constant.

We interpret the eigenstate Ψ as the state of a particle with definite energy, traveling within a region of uniform gravitational field and interacting with a quantum clock. In order to determine the time of flight, we follow the ideas presented in Sec. 6 and suppose that the interaction between the particle and the clock barrier can be treated as a small perturbation. At this point it is interesting to compare the present situation with that of the Sec. 6. There, the scattered (transmitted) wave function could be written as a product of a free particle wave function by a clock state carrying the information about the time of flight. In the gravitational case the situation is analogous, since that in the limit of small perturbations the only effect of the interaction particle-clock is to make the clock run and record the time of flight. In the following, we will show that in the far future the scattered (reflected) wave function can be written as a product of a wave function of a particle under the action of a purely gravitational potential (without the clock) by a clock state showing the time of flight.

We now turn our attention to the reflected wave function in the far future. In the stationary framework that means that we must choose $\xi < 0$ and $|\xi|$ "large". This fact allows us to make use of the asymptotic form of the Airy functions in (90). According to [26], if $\xi < 0$ and $|\xi|$ is sufficiently large we have

$$Ai(\xi) \sim \frac{1}{\sqrt{\pi}} |\xi|^{-1/4} \cos\left(\frac{2}{3}|\xi|^{3/2} - \frac{\pi}{4}\right) \quad , \tag{91}$$

$$Bi(\xi) \sim \frac{1}{\sqrt{\pi}} |\xi|^{-1/4} \sin\left(\frac{2}{3}|\xi|^{3/2} - \frac{\pi}{4}\right) \quad . \tag{92}$$

The trigonometric functions can be written in terms of complex exponentials, and the reflected wave function comes from the exponentials with negative exponents. Using (91) and (92) into (77) we can write the following expression for the reflected wave function

$$\psi^n_{k,ref}(y) \sim \; A\,\frac{1}{\sqrt{\pi}}|\xi|^{-1/4}\,\frac{1}{2}\,e^{-i\left(\frac{2}{3}|\xi|^{3/2}-\frac{\pi}{4}\right)} - B\,\frac{1}{\sqrt{\pi}}|\xi|^{-1/4}\,\frac{1}{2i}\,e^{-i\left(\frac{2}{3}|\xi|^{3/2}-\frac{\pi}{4}\right)} \Longrightarrow$$

$$\Longrightarrow \psi^n_{k,ref}(y) \sim \; A\,\frac{1}{2\sqrt{\pi}}|\xi|^{-1/4}\,\left(1+i\,\frac{B}{A}\right)\,e^{-i\left(\frac{2}{3}|\xi|^{3/2}-\frac{\pi}{4}\right)} \quad . \tag{93}$$

Analogously, the incident wave function reads

$$\psi^n_{k,inc}(y) \sim A\,\frac{1}{2\sqrt{\pi}}|\xi|^{-1/4}\,\left(1-i\,\frac{B}{A}\right)\,e^{i\left(\frac{2}{3}|\xi|^{3/2}-\frac{\pi}{4}\right)} \quad . \tag{94}$$

We now choose A in such a way that $A = 1 + i\,B$. With this choice the function $\psi^n_k(y)$ reduces to $Ai(\frac{y-b}{a})$ in the limit of a negligible clock barrier. As a consequence, the reflected and the incident waves read

$$\psi^n_{k,ref}(y) \sim \left(\frac{1+i\frac{B}{A}}{1-i\frac{B}{A}}\right)\,\frac{1}{2\sqrt{\pi}}|\xi|^{-1/4}\,e^{-i\left(\frac{2}{3}|\xi|^{3/2}-\frac{\pi}{4}\right)} \quad , \tag{95}$$

$$\psi^n_{k,inc}(y) \sim \frac{1}{2\sqrt{\pi}}|\xi|^{-1/4}\,e^{i\left(\frac{2}{3}|\xi|^{3/2}-\frac{\pi}{4}\right)} \quad . \tag{96}$$

Writing the complex constant $R = \left(\frac{1+i\frac{B}{A}}{1-i\frac{B}{A}}\right)$ in the exponential form, *i.e.*, $R = |R|\,e^{i\,\varphi_R} = \; e^{i\,\varphi_R}$, we get

$$\psi^n_{k,ref}(y) \sim \; \frac{1}{2\sqrt{\pi}}|\xi|^{-1/4}\,e^{-i\left(\frac{2}{3}|\xi|^{3/2}-\frac{\pi}{4}-\varphi_R\right)} \quad , \tag{97}$$

where the phase shift φ_R can be caracterized as

$$\varphi_R = \arctan\left(\frac{2\frac{B}{A}}{1-\frac{B^2}{A^2}}\right) \quad . \tag{98}$$

The phase shift φ_R is a function of the energy of the clock barrier ($E_{cl} = n\hbar\omega$). In particular, in the absence of the barrier, we have $\varphi_R(0) = 0$, because if the potential is purely gravitational, the stationary state is the Airy function $Ai(\xi)$, as showed in Sec. 4. For a very low energy barrier, *i.e.*, for $n\hbar\omega << E$, we can expand φ_R in a power series of $n\hbar\omega$ and restrict ourselves to its first nonvanishing term, the first order correction to the phase shift. This term will give rise to the shift in the clock initial state, which will allow us to identify the time of flight of the particle.

We already know the zeroth order term in the power series expansion ($\varphi_R(0) = 0$). Let us now calcule the first order correction. According to (98), we have

$$\varphi_R(n\hbar\omega) = \frac{\partial}{\partial E_{cl}} \arctan\left(\frac{2\frac{B}{A}}{1-\frac{B^2}{A^2}}\right)\Bigg|_{E_{cl}=0} \cdot n\hbar\omega + \cdots \quad . \tag{99}$$

By a direct calculation we get

$$\frac{\partial}{\partial E_{cl}} \arctan\left(\frac{2\frac{B}{A}}{1-\frac{B^2}{A^2}}\right) = \frac{2}{1+\frac{B^2}{A^2}} \frac{\partial}{\partial E_{cl}}\left(\frac{B}{A}\right) \quad . \tag{100}$$

Substituting (100) into (99) and using the fact that $\frac{B}{A}(0) = 0$, we find

$$\varphi_R(n\hbar\omega) = 2\frac{\partial}{\partial E_{cl}}\left(\frac{B}{A}\right)\Bigg|_{E_{cl}=0} \cdot n\hbar\omega + \cdots \quad . \tag{101}$$

According to (85), the derivative of $\frac{B}{A}$ with respect to E_{cl} (calculated at $E_{cl} = 0$) gives

$$\frac{\partial}{\partial E_{cl}}\left(\frac{B}{A}\right)\Bigg|_{E_{cl}=0} = \frac{1}{Bi\,(\xi_0)} \frac{\partial}{\partial E_{cl}}\left[\frac{C}{A} Ai\left(\xi_0 + \frac{E_{cl}}{mga}\right)\right]\Bigg|_{E_{cl}=0} \quad . \tag{102}$$

The derivative of $\frac{C}{A}$ calculated at $E_{cl} = 0$ reads

$$\frac{\partial}{\partial E_{cl}}\left(\frac{C}{A}\right)\Bigg|_{E_{cl}=0} = -\pi\ \frac{\partial Ai}{\partial E_{cl}}\Bigg|_{E_{cl}=0} \frac{dBi}{d\xi}(\xi_0) + \pi\, Bi\ \frac{\partial}{\partial E_{cl}}\left(\frac{dAi}{d\xi}\right)\Bigg|_{E_{cl}=0} \quad , \tag{103}$$

where we have made use of the Wronskian identity (87).

Besides that, we have

$$\frac{\partial Ai}{\partial E_{cl}}\Bigg|_{E_{cl}=0} = \frac{1}{mga}\frac{dAi}{d\xi}(\xi_0) \quad \text{and} \quad \frac{\partial}{\partial E_{cl}}\frac{dAi}{d\xi}\Bigg|_{E_{cl}=0} = \frac{1}{mga}\frac{d^2Ai}{d\xi^2}(\xi_0) \quad . \tag{104}$$

By firstly substituting (104) into (103) and then coming back to (102) we get

$$\frac{\partial}{\partial E_{cl}}\left(\frac{B}{A}\right)\Bigg|_{E_{cl}=0} = \frac{1}{mga\,Bi}\left[-\pi\,\frac{dAi}{d\xi}\frac{dBi}{d\xi}Ai + \pi\,Bi\frac{d^2Ai}{d\xi^2}Ai + \frac{dAi}{d\xi}\right] \quad . \tag{105}$$

We note that all functions in the right hand side of (105) are calculated at ξ_0.

Making use of the Wronskian identity (87) again we can write

$$\frac{\partial}{\partial E_{cl}} \left(\frac{B}{A}\right)\bigg|_{E_{cl}=0} = \frac{\pi}{mga} \left[Ai \frac{d^2 Ai}{d\xi^2} - \frac{dAi}{d\xi} \frac{dAi}{d\xi}\right] \quad . \tag{106}$$

Now, by using the Airy equation into (106) we get

$$\frac{\partial}{\partial E_{cl}} \left(\frac{B}{A}\right)\bigg|_{E_{cl}=0} = \frac{\pi}{mga} \left\{\xi_0 \left[Ai(\xi_0)\right]^2 - \left[\frac{dAi}{d\xi}(\xi_0)\right]^2\right\} \quad . \tag{107}$$

Since the argument ξ_0 is a negative number of large absolute value, we can use the asymptotic form of the functions $Ai(\xi)$ and $\frac{dAi}{d\xi}$. We already know the former (see Eq. (91)). The later can be found in [26]. It reads (up to its leading term)

$$\frac{dAi}{d\xi} \sim \frac{1}{\sqrt{\pi}} |\xi|^{1/4} \sin\left(\frac{2}{3}|\xi|^{3/2} - \frac{\pi}{4}\right) \quad . \tag{108}$$

Using (91) and (108) into (107) we find $\frac{\partial}{\partial E_{cl}} \left(\frac{B}{A}\right)\big|_{E_{cl}=0} = -\frac{1}{\hbar}\sqrt{2\left(\frac{b-y_0}{g}\right)}$, so that (101) can be written as

$$\varphi_R(n\hbar\omega) = -2n\omega\sqrt{\frac{2\,(b-y_0)}{g}} + \cdots \quad . \tag{109}$$

We now write $T \equiv 2\sqrt{\frac{2(b-y_0)}{g}}$ and substitute (109) into (97) to get the explicit expression of $\psi^n_{k,ref}(y)$

$$\psi^n_{k,ref}(y) \sim \frac{1}{2\sqrt{\pi}} |\xi|^{-1/4} \, e^{-i\left(\frac{2}{3}|\xi|^{3/2} - \frac{\pi}{4}\right)} \, e^{-in\omega T} \quad . \tag{110}$$

The reflected part of the eigenstate $\Psi(x, y, q)$ (see Eq. (89)) reads

$$\begin{aligned} \Psi_{E,k,ref}(x, y, q) &= \mathcal{C}_{E,k} \frac{e^{ikx}}{2\sqrt{\pi}} |\xi|^{-1/4} \, e^{-i\left(\frac{2}{3}|\xi|^{3/2} - \frac{\pi}{4}\right)} \sum_{n=-j}^{j} \frac{e^{in(q-\omega T)}}{\sqrt{2\pi N}} \qquad (111) \\ &= \mathcal{C}_{E,k} \frac{e^{ikx}}{2\sqrt{\pi}} |\xi|^{-1/4} \, e^{-i\left(\frac{2}{3}|\xi|^{3/2} - \frac{\pi}{4}\right)} \, v_0(q - \omega T) \quad . \end{aligned}$$

Following the ideas of Sec. 6 (see Eq. (59) and the discussion below that equation) we recognize the shifted clock state $v_0(q-\omega T)$ as one of the factors in the reflected wave function, as expected. We can thus finally say that the time of flight for a quantum particle in a uniform gravitational field, in the context of NCQM, is given by

$$T = 2\sqrt{\frac{2\,(b-y_0)}{g}} \quad . \tag{112}$$

We note that (112) is equal to the corresponding expression of the ordinary quantum mechanics [23]. More than this, we see that the time of flight in NCQM is equal to the classical one.

The main result of this section is Eq. (112), which corresponds, in the sense we have been discussing in this work, to an extention of the Equivalence Principle to the realm of NCQM.

To finish, it is important to clarify the differences between the approach we have used (Peres approach [25]) in computing the time of flight of a particle in a uniform gravitational field (in NCQM) and that by Davies (in ordinary quantum mechanics) [23].

First of all, it is important to bear in mind that the central idea of Peres is to couple a particle to a quantum system (that plays the role of a quantum rotor), which will work as a quantum clock. When this coupling is considered small, it is such that the effect of the particle-clock interaction is essentially only to cause a displacement of the (initial) position of the clock pointer (*i.e.*, to change the initial state of the clock). Thus, at the end of the interaction, the clock will record the time of flight of the particle, which is simply the time interval during which the particle interacted with the clock-rotor.

In order that one can "watch" the final position of the clock pointer, and thus get the time of flight of the particle, it is necessary that a measurement be realized. This, in its turn, must be done in the "distant future", when the particle-clock interaction can be considered totally negligible. That is one of the key points of Peres approach. By analysing the scattered wave function, we have thus shown that the time of flight is codified in its phase shift (see Eq.'s (109)-(112)). This result is totally in accordance with the spirit of that proved by Peres in the case of a free particle [25].

In the approach by Davies for computing the time of flight of a particle in a uniform gravitational field, the Hamiltonian of the system does not contain the coupling term between the particle and the clock. In this case, the solution of the scattering problem do not "see" the clock, but only the gravitational barrier. As such, what Davies determined is not the phase shift of the wave function due to the interaction between the particle (under the action of gravity) and the clock barrier. In fact, what he has done was to calculate the derivative (with respect to the total energy) of the phase difference induced by the action solely of gravity on the particle wave function. This derivative was calculated at the point where the clock detector should have been placed (it should be noted that this derivative is position dependent). Nevertheless, it follows that the final result for the time of flight obtained by Davies turns out to be the same that would be obtained by Peres approach, described above. Hence, in Davies approach the clock is introduced *ad hoc* for the computation of the time of flight, through the prescription that the derivative of the phase difference of the particle wave function has to be calculated at the point which corresponds to the position where the clock detector should be placed. We remark that in Peres approach, which is based on the calculation of the phase shift of the wave function, there is no need of dealing with position dependent phases.

In our application of computing the time of flight we have chosen to explicitly perform the calculations of the scattering theory. As a by product, when we take the non-commutativity parameter equal to zero, we obtain the extension of the Peres result for the time of flight of a particle to the case that, instead of being free, it is under the action of a uniform gravitational field.

8 Concluding remarks

We have studied the gravitational quantum well in non-commutative quantum mechanics. Assuming non-commutativity only on configuration space, we have *exactly* solved the non-commutative Schrödinger equation. As in ordinary quantum mechanics, in the non-commutative case the solution is given by a Airy function and the energy eigenvalues are expressed in terms of the zeroes of the Airy function Ai. Then, from data of the quantum gravitational well experiment with freely falling neutrons (the Grenoble experiment) [16]-[18] we could then set an upper-bound on the value of the spatial non-commutative parameter, θ. This result can be improved in the future, when more accurate experimental data will be available [34]. We note that, as far as we now, the works that considered the non-commutative gravitational quantum well have not established an upper-bound on θ, but just on the momentum-momentum non-commutativity parameter, as they investigated the problem by assuming non-commutativity on phase space, as well as approached it perturbatively.

Another issue we have addressed in this paper was related to the question of the (Weak) Equivalence Principle, its validity in NCQM. We were interested in investigating the status of the Equivalence Principle through a kind of (balistic) Galileo experiment, associated to delocalized, energy eigenstates. For that, we have used a quantum clock model, due to Peres [25], in order the time of flight of the particle could be measured. We remark that although we were inspired by Davies [23] in asking for a possible violation of the Equivalence Principle by studying the time of flight of a particle subjected to a uniform gravitational field, we have performed a conventional stationary state analysis of a scattering problem, instead of just naively applying the formula used by Davies in the ordinary case [23]. Instead of this, we closely followed the original approach by Peres [25] of applying a quantum clock in the measurement of the time of flight of a quantum particle. It resulted that the time of flight is the same as in quantum mechanics, which in turn, is identical to the classical result, when the measurement is made far from the turning point. This fact suggests that it might be possible that one has an extention of the (Weak) Equivalence Principle to NCQM.

In order to address until to what extent the (Weak) Equivalence Principle holds in NCQM (and analogously in ordinary quantum mechanics) further studies are important, such as, for instance, the investigation of how matter and anti-matter, in a non-commutative background, behave under the action of a gravitational field. Although it has been shown that in NCQFTs the CPT symmetry is still preserved [35], but with indidual violations of the C, T and P symmetries, such a symmetry might be broken at a more fundamental level, when not only the matter fields are quantized but also the gravitational field itself. In fact, it has been suggested that quantum gravity effects might lead to violations of the CPT symmetry [36]. Finally, we recall that as the non-commutativity of space-time might be a signature of quantum gravity, experiments involving an interplay between gravity and quantum mechanics are welcome, since even at low-energy, but with enough level of accuracy, they might display information about the physics whose origin is in a more fundamental level (see [37]).

It would be interesting to experimentally investigate the Equivalence Principle by measuring the time of flight of quantum particles using a quantum clock. Of course that one has to deal with the intrinsic uncertainty of using a quantum clock, as even at low energy (weak coupling between the clock and the system of interest), a good measurement of the time of flight is subjected to a lower limit on the time resolution of the clock [25]. In this direction, we quote the work of Alonso *et al.* [38], which has shown, in the case of a free quantum particle, that it is possible to gain an improvement in the measurement of time of flight if the quantum clock does not work continuouly but rather by means of pulsed couplings.

Acknowledgments

The authors would like to dedicate this work *in memoriam* to *Ivens Carneiro*, a great friend and admirable physicist, who prematurely passed away. K. H. C. C-B acknowledges Prof. D. A. T. Vanzella, for encouragement, and full support from the *Fundação de Amparo à Pesquisa no Estado de São Paulo* (FAPESP). A. G. Martins acknowledges the *Universidade de Brasília* (UnB), for support during the writing of this paper.

References

[1] *Wolfgang Pauli, Scientific Correspondence*, Vol II, p.15, Ed. Karl von Meyenn, Springer-Verlag, 1985; *ibidem*, Vol. III, p.380, 1993.

[2] H. S. Snyder, *Phys. Rev.* **71**, 38 (1947).

[3] R. J. Szabo, *Phys. Rept.* **378**, 207 (2003).

[4] R. Jackiw, <http://arxiv.org/abs/hep-th/0110057>.

[5] A. Connes, M. Douglas, and A. S. Schwarz, *J. High Energy Phys.* **02**, 003 (1998); N. Seiberg and E. Witten, *J. High Energy Phys.* **09**, 032 (1999).

[6] M. Douglas and N. A. Nekrasov, *Rev. Mod. Phys.* **73**, 977 (2001).

[7] H. O. Girotti, <http://arxiv.org/abs/hep-th/0301237>.

[8] S. Doplicher, K. Fredenhagen, and J. E. Roberts, *Phys. Lett.* **B331**, 39 (1994); *ibidem*, *Commun. Math. Phys.*, **172**, 187 (1995).

[9] L. Mezincescu, *hep-th/0007046*; J. Gamboa, M. Loewe, and J. C. Rojas, *Phys. Rev.* **D64**, 067901 (2001); V. P. Nair and A. P. Polychronakos, *Phys. Lett.* **B505**, 267 (2001); B. Morariu and A. P. Polychronakos, *Nucl. Phys.* **B610**, 531 (2001). S. Bellucci, A. Neressian and C. Sochichiu, *Phys. Lett.* **B522**, 345 (2001); H. R. Christiansen and F. A. Schaposnik, *Phys. Rev.* **D65**, 086005 (2002); J. Gamboa, M. Loewe, F. Mendez, and J. C. Rojas, *Int. J. Mod. Phys.* **A17**, 2555 (2002); S. Bellucci and A. Yeranyan, *Phys. Rev.* **D72**, 085010 (2005); S. Bellucci and A. Yeranyan, *Phys. Lett.* **B609**,418 (2005); X. Calmet and M. Selvaggi, *Phys. Rev.* **D74**, 037901 (2006).

[10] J.-Zu. Zhang, *Phys. Rev.* **D74**, 124005 (2006); Y. Wu and X. Yang, *Phys. Rev.* **D73**, 067701 (2006); J.-Zu. Zhang, *Phys. Rev. Lett.* **93**, 043002 (2004); Z. Guralnik, R. Jackiw, S. Y. Pi, and A. P. Polychronakos, *Phys. Lett.* **B517**, 450 (2001); J. Gamboa, M. Loewe, and J. C. Rojas, hep-th/0101081.

[11] Y.-H. Luo and Z.-m. Ge, *Commun. Theor. Phys.* **46**, 968 (2006); F. S. Bemfica and H. O. Girotti, *J. Phys.* **A38**, L539 (2005); P.-M. Ho and H.-C. Kao, *Phys. Rev. Lett.* **88**, 151602 (2002).

[12] A. P. Balachandran, T. R. Govindarajan, C. Molina, and P. Teotonio-Sobrinho, *JHEP* **0410**, 072 (2004).

[13] A. P. Balachandran, T. R. Govindarajan, A. G. Martins, and P. Teotonio-Sobrinho. *JHEP* **0411**, 068 (2004).

[14] A. P. Balachandran, A. G. Martins and P. Teotonio-Sobrinho, *JHEP* **0705**, 066 (2007).

[15] P. W. Langhoff, *Am. J. Phys.* **39**, 954 (1971); J. Gea-Banacloche, *Am. J. Phys.* **67**, 776 (1999); O. Vallee, *Am. J. Phys.* **68**, 672 (2000); M. A. Doncheski and R. W. Robinett, *Am. J. Phys.* **69**, 1084 (2001); S. Flügge, *Practical Quantum Mechanics I* (Berlin, Springer, 1974); L. D. Landau and E. M. Lifshitz, *Quantum Mechanics*, 3rd. Edition, Pergamon Press, Oxford (1976); J. J. Sakurai, *Modern Quantum Mechanics*. Prentice-Hall, 1994.

[16] V. V. Nesvizhevsky *et al.*, *Nature* **415**, 297 (2002).

[17] V. V. Nesvizhevsky *et al.*, *Phys. Rev.* **D67**, 102002 (2003).

[18] V. V. Nesvizhevsky *et al.*, *Eur. Phys. J.* **C40**, 479 (2005).

[19] O. Bertolami, J. G. Rosa, C. M. L. de Aragão, P. Castorina, and D. Zappala, *Phys. Rev.* **D72**, 025010 (2005).

[20] R. Banerjee, B. D. Roy, and S. Samanta. *Phys. Rev.* **D74**, 045015 (2006).

[21] A. Saha, *Eur. Phys. J.* **C51**, 199 (2007).

[22] F. Buisseret, V. Mathieu, and B. Silvestre-Brac, *Class. Quant. Grav.* **24**, 855 (2007); F. Brau and F. Buisseret, *Phys. Rev.* **D74**, 036002 (2006).

[23] P. C. W. Davies, *Class. Quantum Grav.*, **21**, 2761 (2004).

[24] O. Bertolami, J. G. Rosa, C. M. L. de Aragão, P. Castorina, and D. Zappalà, *Mod. Phys. Lett.* **A21**, 795 (2006).

[25] A. Peres, *Am. J. Phys.*, **48**, 552 (1980).

[26] F. W. J. Olver, *Asymptotics and Special Functions.* New York: Academic Press, 1974.

[27] R. Colella, A. W. Overhauser, and S. A. Werner, *Phys. Rev. Lett.*, **34**, 1472 (1975).

[28] S. M. Carroll, J. A. Harvey, V. Alan Kostelecky, C. D. Lane, and T. Okamoto, *Phys. Rev. Lett.* **87**, 141601 (2001).

[29] M. Chaichian, M. M. Sheikh-Jabbari, and A. Tureanu, *Phys. Rev. Lett.* **86**, 2716 (2001).

[30] O. Bertolami and F. M. Nunes, *Class. Quantum Grav.* **20**, L61 (2003).

[31] A. Peres, *Quantum Theory: Concepts and Methods.* Kluwer Academic Publishers. Dordrecht, The Netherlands (1993).

[32] C. R. Leavens, *Solid State Commun.*, **86**, 781 (1993).

[33] L. Viola and R. Onofrio, *Phys. Rev.* **D55**, 455 (1997).

[34] V. V. Nesvizhevsky *et al.*, *Eur. Phys. J.* **C40**, 479 (2005).

[35] M. Chaichian, K. Nishijima, and A. Tureanu, *Phys. Lett.* **B568**, 146 (2003).

[36] S. W. Hawking, *Phys. Rev.* **D14**, 2460 (1976); R. M. Wald, *Phys. Rev.* **D21**, 2742 (1980).

[37] G. Amelino-Camelia, *Phys. Rev.* **D62**, 024015 (2000).

[38] D. Alonso, R. Sala Mayato, and J. G. Muga, *Phys. Rev.* **A67**, 032105 (2003).